

SEVENTH FRAMEWORK PROGRAMME  
THEME 6: Environment (including climate change)




Contract for:

Collaborative Project

### *D.3.2.3*

## *GIS data maps of energy and water fluxes, pollution concentrations, land cover and vegetation*



Project acronym: BRIDGE  
Project full title: sustainaBle uRban planning Decision support accountinG for urban mEtabolism  
Contract no.: 211345  
Date: 18/05/2011  
Document Reference: 211345\_014\_TR\_CMCC  
Book Captain: Donatella Spano (CMCC)  
Authors: Donatella Spano (CMCC)  
Serena Marras (CMCC)  
Contributors: Beniamino Gioli (CNR-IBIMET)  
Piero Toscano (CNR-IBIMET)  
Sue Grimmond (KCL)  
M. J. Tallis (SOTON)  
Timo Vesala (UHEL)  
Tomasz Staszewski (IETU)  
Stathopoulou Marina (NKUA)  
Nektarios Chrysoulakis (FORTH)  
Zina Mitraka (FORTH)  
Issue: 1.0  
Deliverable no.: D.3.2.3  
Dissemination Level: CO



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 2/32

## Document Status Sheet

Issue	Date	Author	Comments
1.0	18/5/2011	ENZO MAGLIULO (CNR)	UPDATES FROM CMCC.



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
 Contract no.: 211345  
 Document Ref.: 211345\_014\_TR\_CMCC  
 Issue: 1.0  
 Date: 18/05/2011  
 Page number: 3/32

## Table of Contents

<b>1. INTRODUCTION .....</b>	<b>4</b>
1.1 Purpose of the document .....	4
1.2 Definitions and Acronyms .....	4
1.3 Document References .....	4
1.4 Project Overview .....	5
<b>2. SUMMARY .....</b>	<b>7</b>
2.1 Partners involved .....	7
<b>3. CASE STUDIES.....</b>	<b>9</b>
3.1 Helsinki .....	9
Data already existing .....	9
Data collected during the project .....	12
Contact info .....	12
3.2 Athens.....	12
Data already existing .....	12
Data collected during the project .....	18
Contact info .....	18
3.3 London .....	20
Data already existing .....	20
Data to be collected during the project .....	21
Contact info .....	23
3.4 Firenze .....	24
Data already existing .....	24
Data collected during the project .....	24
Contact info .....	24
3.5 Gliwice .....	26
Data already existing .....	26
Data collected during the project .....	26
Contact info .....	26
<b>ANNEX:.....</b>	<b>27</b>



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 4/32

## 1. Introduction

The city is a complex system that, in order to maintain its vital functions, consumes materials and energy, accumulates materials and releases different forms of waste. In the BRIDGE (sustainaBle uRban plannIng Decision support accountinG for urban mEtabolism) project, the concept of “*urban metabolism*” is used to indicate the exchange and transformation of energy and matter between a city and its environment. Urban surface structures, such as buildings, significantly influence the **water balance**, the **energy balance** and **air quality**. In addition, land use change due to construction of buildings, roads and factories, determine major changes at the regional scale. Exchanges of energy, heat, moisture, carbon and pollutant are then measured and modelled, during the project, using recent advances in urban climatology.

The role of WP3 is to collect and analyze valuable datasets and to describe over time methodologies used to estimate urban exchanges. Five European cities have been selected (Helsinki, Athens, London, Firenze and Gliwice) as representative of the different city typologies and influenced by different policy and resource availability. For each case study, direct measurements will be performed in order to estimate urban exchanges and models will be used to simulate and predict these fluxes. In addition, the role of land use in relation to urban pattern and typology will be assessed using remote sensing data and methodologies.

### 1.1 Purpose of the document

This document is the D.3.2\_Remote Sensing Data Collection and Analysis. Task 3.2 has the role to systematically monitor the main fluxes using remote sensing techniques for each case study (Helsinki, Athens, London, Firenze and Gliwice).

The **aim of this document** is to collect information on remote sensing data available for each case study of the BRIDGE Project. In particular, it contains information on data spatio-temporal distributions, methodologies used to provide the data and maps of physical parameters. This information will be useful to know the state-of-art in remote sensing studies and data for each city involved. The final products of Task 3.2, in fact, are GIS data and maps of energy and water fluxes, pollution concentrations, land cover and vegetation.

### The present document updates D3.2.2

### 1.2 Definitions and Acronyms

#### Acronyms

CoP	Community of Practice
DSS	Decision Support System
GIS	Geographical Information System
RS	Remote Sensing
DEM	Digital Elevation Model
ASTER	Advanced Thermal Emission and Reflection Radiometer

### 1.3 Document References

- Chrysoulakis, N., Abrams, M., Kamarianakis, Y. and Stanisławski, m., 2009. Validation of the ASTER derived Global DEM product (GDEM) for the area of Greece. ISPRS Journal of Photogrammetry and Remote Sensing (accepted).
- Falk M., Pyles R.D., Marras S., Spano D., Snyder R.L., Paw U K.T. Coupling of the WRF and ACASA models for urban environments: two case studies. AMS Conference on Ninth Symposium on the Urban Environment, Keystone, Colorado, USA, August 2-6, 2010.



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.:	D.3.2.3
Contract no.:	211345
Document Ref.:	211345_014_TR_CMCC
Issue:	1.0
Date:	18/05/2011
Page number:	5/32

- Fuller, R.M., Smith, G.M., Sanderson, J.M., Hill, R.A., & Thomson, A.G. 2001. The UK Land Cover Map 2000: construction of a parcel-based vector map from satellite images. *Cartographic J.* **39** 15-25.
- Jiménez-Munõz, J.C. and Sobrino, J.A., 2003. A generalized single-channel method for retrieving land surface temperature. *Journal of Geophysical Research*, 108 (D22), 4688, doi:10.1029/2003JD003480.
- LAEI 2006. London Atmospheric Emissions Inventory 2006. Compiled by AEA, March 2009. [http://static.london.gov.uk/mayor/environment/air\\_quality/research/emissions-inventory.jsp](http://static.london.gov.uk/mayor/environment/air_quality/research/emissions-inventory.jsp)
- Liang, S., 2001. Narrowband to Broadband Conversion of Land Surface Albedo. I. Algorithms, *Remote Sensing of Environment*, vol. 76, pp. 213-238.
- Liang, S., 2004. *Quantitative Remote Sensing of Land Surfaces*, John Wiley and Sons, Inc., 534 pages.
- Nowak, D.J. 1994. Air pollution removal by Chicago's Urban Forest. In: McPherson E.G., Nowak D.J., Rowntree R.A. (Eds). *Chicago's Urban Forest Ecosystem: Results of the Chicago Urban Forest Climate Project*. US Department of Agriculture, Forest Service, Northeastern Forest Experiment Station, Radnor, PA, pp. 63-82.
- Sobrino, J.A., Jimenez-Munoz, J.C. and Paolini, L., 2004. Land surface temperature retrieval from LANDSAT TM 5. *Remote Sensing of Environment*, vol. 90, pp. 434-440.
- Stathopoulou, M., Cartalis, C. and Petrakis, M., 2007. Integrating Corine Land Cover data and Landsat TM for surface emissivity definition: application to the urban area of Athens, Greece. *International Journal of Remote Sensing*, vol. 28(15), pp. 3291 – 3304.

## 1.4 Project Overview

Urban metabolism considers a city as a system and distinguishes between energy and material flows. “Metabolic” studies are usually top-down approaches that assess the inputs and outputs of food, water, energy, etc. from a city, or that compare the metabolic process of several cities. In contrast, bottom-up approaches are based on quantitative estimates of urban metabolism components at local scale, considering the urban metabolism as the 3D exchange and transformation of energy and matter between a city and its environment. Recent advances in biophysical sciences have led to new methods to estimate energy, water, carbon and pollutants fluxes. However, there is poor communication of new knowledge to end-users, such as planners, architects and engineers.

BRIDGE aims at illustrating the advantages of considering environmental issues in urban planning. BRIDGE will not perform a complete life cycle analysis or whole system urban metabolism, but rather focuses on specific metabolism components (energy, water, carbon, pollutants). BRIDGE's main goal is to develop a Decision Support System (DSS) which has the potential to propose modifications on the metabolism of urban systems towards sustainability.

BRIDGE is a joint effort of 14 Organizations from 11 EU countries. Helsinki, Athens, London, Firenze and Gliwice have been selected as case study cities. The project uses a “Community of Practice” approach, which means that local stakeholders and scientists of the BRIDGE meet on a regular basis to learn from each other. The end-users are therefore involved in the project from the beginning. The energy and water fluxes are measured and modelled at local scale. The fluxes of carbon and pollutants are modelled and their spatio-temporal distributions are estimated. These fluxes are simulated in a 3D context and also dynamically by using state-of-the-art numerical models, which normally simulate the complexity of the urban dynamical process exploiting the power and capabilities of modern computer platforms. The output of the above models lead to



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.:	D.3.2.3
Contract no.:	211345
Document Ref.:	211345_014_TR_CMCC
Issue:	1.0
Date:	18/05/2011
Page number:	6/32

indicators which define the state of the urban environment. The end-users decide on the objectives that correspond to their needs and determine objectives' relative importance. Once the objectives have been determined, a set of associated criteria are developed to link the objectives with the indicators. BRIDGE integrate key environmental and socio-economic considerations into urban planning through Strategic Environmental Assessment. The BRIDGE DSS evaluates how planning alternatives can modify the physical flows of the above urban metabolism components. A Multi-criteria Decision Making approach has been adopted in BRIDGE DSS. To cope with the complexity of urban metabolism issues, the objectives measure the intensity of the interactions among the different elements in the system and its environment. The objectives are related to the fluxes of energy, water, carbon and pollutants in the case studies. The evaluation of the performance of each alternative is done in accordance with the developed scales for each criterion to measure the performance of individual alternatives.

Several studies have addressed urban metabolism issues, but few have integrated the development of numerical tools and methodologies for the analysis of fluxes between a city and its environment with its validation and application in terms of future development alternatives, based on environmental and socio-economic indicators for baseline and extreme situations. The innovation of BRIDGE lies in the development of a DSS integrating the bio-physical observations with socio-economic issues. It allows end-users to evaluate several urban planning alternatives based on their initial identification of planning objectives. In this way, sustainable planning strategies will be proposed based on quantitative assessments of energy, water, carbon and pollutants fluxes.



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 7/32

## 2. Summary

Task 3.2 aims to collect satellite, airborne and ground-based remote sensing data in order to produce maps of energy and water fluxes, pollution concentration, land cover and vegetation.

For each case study, the following data will be available:

- Maps of surface albedo in the visible part of the spectrum (NKUA) by satellite data
- Urban land cover/land use maps (NKUA, SOTON, UHEL)
- Land surface temperature (NKUA, KCL) by satellite data
- Morphological characteristics and anthropogenic heat flux (KCL)
- Maps of sensible heat flux based on different modelling approaches (KCL, CMCC)
- Maps of surface emissivity (CMCC, NKUA) by modelling and satellite data
- 3D maps of cities
- Particulate profiles and PBL characteristics (CNR) by LIDAR
- Airborne images (UHEL, KCL, CNR)
- Digital Elevation Models (FORTH)
- High resolution satellite data products (FORTH)

More details on data spatial and temporal variability and methodologies used are in the following sections.

### 2.1 Partners involved

#### 1. CNR

Consiglio Nazionale delle Ricerche, Italy

Enzo Magliulo ([enzo.magliulo@cnr.it](mailto:enzo.magliulo@cnr.it))

Franco Miglietta ([f.miglietta@ibimet.cnr.it](mailto:f.miglietta@ibimet.cnr.it)): Firenze case study leader

#### 2. CMCC

Centro Euro-Mediterraneo per i Cambiamenti Climatici S.c.a.r.l, Italy

Donatella Spano ([spano@uniss.it](mailto:spano@uniss.it))

#### 3. KCL

King's College London

Sue Grimmond ([sue.grimmond@kcl.ac.uk](mailto:sue.grimmond@kcl.ac.uk)): London case study leader

#### 4. NKUA

National and Kapodistrian University of Athens, Greece

Mattheos Santamouris ([msantam@phys.uoa.gr](mailto:msantam@phys.uoa.gr)): Athens case study leader

#### 5. FORTH

Foundation for Research and Technology – Hellas, Greece

Nektarios Crysoulakis ([zedd2@iacm.forth.gr](mailto:zedd2@iacm.forth.gr))

#### 6. SOTON

School of Biological Sciences-United Kingdom

Prof. Gail Taylor ([G.Taylor@soton.ac.uk](mailto:G.Taylor@soton.ac.uk))



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 8/32

---

### 7. UHEL

University of Helsinki, Finland

Timo Vesala ([timo.vesala@helsinki.fi](mailto:timo.vesala@helsinki.fi)): Helsinki case study leader

### 8. IETU

Instytut Ekologii Terenów Uprzemysłowionych, Poland

Tomasz Staszewski ([stasz@ietu.katowice.pl](mailto:stasz@ietu.katowice.pl)): Gliwice case study leader




## 3. Case Studies

### 3.1 Helsinki




#### Data already existing

- *GIS data.* Examples are given in Figure 1.
- *Aerial images*



**Figure 1.** Helsinki Land Use Map.

**Table 1.** Land Use Classification in Slices. Raster image in 10 m x 10 m pixels, coordinate system EUREF-FIN/TM35FIN = ETRS89, Transverse Mercator, Zone 35.

Land use class	Colour
<b>A. Housing and recreational areas</b>	(main class)
<b>A1. Housing areas</b>	(main class)
<b>A11. Block house areas</b>	
<b>A12. Small house areas</b>	(main class)
<b>A121. Row and attached small houses</b>	
<b>A122. Detached housing areas</b>	
<b>A2. Holiday and travel sites</b>	(main class)



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
 Contract no.: 211345  
 Document Ref.: 211345\_014\_TR\_CMCC  
 Issue: 1.0  
 Date: 18/05/2011  
 Page number: 10/32

Land use class	Colour
<b>A21. Holiday resorts and cottage areas</b>	
<b>A22. Areas of travel services and holidaymaking</b>	(main class)
<b>A221. Camping and caravan sites</b>	(no data)
<b>A222. Allotments and cultivation plots</b>	(no data)
<b>A3. Other recreational areas</b>	(main class)
<b>A31. Areas of entertainment services</b>	(no data)
<b>A32. Areas of sport and recreational services</b>	
<b>A33. Parks</b>	
<b>B. Business, management and industrial areas</b>	(main class)
<b>B1. Areas of business and management</b>	(main class)
<b>B11. Commercial and office buildings</b>	(main class)
<b>B111. Commercial buildings</b>	
<b>B112. . Office buildings</b>	
<b>B12. Public buildings</b>	
<b>B2. Industrial and storage areas</b>	(main class)
<b>B21. Industrial areas</b>	
<b>B22. Storage areas</b>	(main class)
<b>B221. Storage buildings</b>	
<b>B222. Other storage areas</b>	(no data)
<b>C. Areas of supporting functions</b>	(main class)
<b>C1. Transportation areas</b>	
<b>C11. Road transportation areas</b>	
<b>C111. Public roads</b>	
<b>C112. Streets and alleys (incl. in C11)</b>	(no data)
<b>C113. Private roads (incl. in C11)</b>	(no data)
<b>C12. Railway areas</b>	
<b>C13. Areas of air traffic and aviation</b>	
<b>C14. Port areas</b>	
<b>C15. Other transportation areas</b>	
<b>C2. Areas of civil engineering</b>	
<b>C21. Environmental engineering and management</b>	
<b>C22. Energy management</b>	
<b>C23. Water management</b>	(no data)
<b>C24. Other sites of civil engineering &amp; management</b>	
<b>D. Rock and soil extraction sites</b>	(main class)
<b>D1. Rock and soil extraction sites</b>	(main class)



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
 Contract no.: 211345  
 Document Ref.: 211345\_014\_TR\_CMCC  
 Issue: 1.0  
 Date: 18/05/2011  
 Page number: 11/32



Land use class	Colour
<b>D11. Rock and mineral extraction sites</b>	(main class)
<b>D111. Mines</b>	
<b>D112. Quarries</b>	
<b>D12. Soil extraction sites</b>	(main class)
<b>D121. Peat production sites</b>	
<b>D122. Gravel and sand extraction sites</b>	
<b>D123. Other soil extraction sites</b>	
<b>E. Agricultural areas</b>	(main class)
<b>E1. Agricultural land in production use</b>	(main class)
<b>E11. Fields</b>	
<b>E12. Perennial meads and grasslands</b>	
<b>E13. Perennial and covered plantations</b>	(main class)
<b>E131. Fruit and berry plantations</b>	
<b>E132. Nursery gardens and covered cultivations</b>	
<b>E2. Other agricultural land</b>	(main class)
<b>E21. Unused agricultural land</b>	(main class)
<b>E211. Long-term fallows</b>	
<b>E212. Agricultural land permanently removed from productive use</b>	
<b>E22. Constructed agricultural land</b>	(main class)
<b>E221. Farm houses and outbuildings</b>	
<b>E222. Other constructed land related to agriculture</b>	
<b>F. Forest management areas</b>	(main class)
<b>F1. Productive forestry areas</b>	(main class)
<b>F11. Forests</b>	
<b>F12. Scrubland</b>	
<b>G. Other land areas</b>	(main class)
<b>G1. Brownfields</b>	
<b>G2. Other land areas</b>	
<b>H. Territorial waters</b>	(main class)
<b>H1. Inland water areas</b>	(main class)
<b>H11. Natural waters</b>	(main class)
<b>H111. Unregulated natural waters</b>	
<b>H112. Regulated natural waters</b>	
<b>H12. Other territorial waters</b>	(main class)
<b>H121. Unregulated waters</b>	



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 12/32

Land use class	Colour
H122. Regulated waters	
H2. Sea areas	

### Data collected during the project

- *3D datasets* can be possibly collected .
- *GIS data*, if more data with the detailed specifications are needed, they can be provided. Maps of sensible heat flux and surface emissivity will be produced by CMCC using the WRF-ACASA model. High resolution maps (200 m) of surface emissivity, sensible and latent heat flux have been produced for the period January-April 2008 (Falk et al., 2010).
- 
- *Satellite data (not via us)*
- *Airborne acquired images*, if more data with the detailed specifications are needed, they can be provided.
- *Digital Elevation Model (DEM)*: Extraction and post processed ASTER GDEM for the broader area of Helsinki was provided by FORTH. The product was delivered in both Geographic WGS84 and UTM WGS84 projections with EGM96 vertical datum. Its spatial resolution is 30 m.

### Contact info

Outi Salminen ([outi.m.salminen@helsinki.fi](mailto:outi.m.salminen@helsinki.fi))  
Annika Nordbo ([annika.nordbo@helsinki.fi](mailto:annika.nordbo@helsinki.fi))


CMCC:  
Prof. Donatella Spano ([spano@uniss.it](mailto:spano@uniss.it))  
Via de Nicola 9, 07100, Sassari, Italy  
Tel. +39 079 229339

Dr. Serena Marras ([serenam@uniss.it](mailto:serenam@uniss.it))  
Via de Nicola 9, 07100, Sassari, Italy  
Tel. +39 079 229372

## 3.2 Athens

### Data already existing

- *3D Maps*
- *GIS data*
  1. Vector data map of Greece used to define the Prefectures of Greece as well as the geographical boundaries of the Attica Prefecture in which Athens is located (Figure 2). Projection info: Lambert Azimuthal Equal-area, map units: meters.



**Figure 2.** Administrative boundaries for Greece.

2. Corine Land Cover 2000 database of Greece: Vector data map at the scale of 1: 100 000, used to describe land cover and land use types across the Prefecture of Attica (including Athens and the Egaleo test site) for the reference year 2000 (Figure 3). The database uses 44 land cover/use classes organized into 5 main categories: 1. Artificial surfaces 2. Agricultural areas 3. Forests and semi-natural areas 4. Wetlands and 5. Water bodies. Projection info: GCS\_ETRS\_1989, map units: degrees





Figure 3. CORINE Land Cover Data for Greece.


- *Aerial images*
- *Satellite data*

On a local scale perspective, processing of the satellite data in the visible part of the spectrum can be performed leading to the measurement and mapping of the following geophysical and biophysical variables for Athens (including the Egaleo test site):

1. Narrowband (spectral) and broadband *surface albedo* at 30m spatial resolution produced from the Landsat TM sensor. The wavelengths (0.25-5.0  $\mu\text{m}$ ) correspond to total shortwave broadband albedo, whereas (0.4-0.7  $\mu\text{m}$ ) and (0.7-5.0  $\mu\text{m}$ ) correspond to visible and near-infrared broadband albedos (Figures 4, 5 and 6). The methodology applied is described in Liang et al. (2002) and Liang (2004).



**Figure 4.** Example image of total shortwave albedo for Athens on 8/4/2005.



**Figure 5.** Example image of visible albedo for Athens on 8/4/2005.




Figure 6. Example image of near-infrared albedo for Athens on 8/4/2005.

2. *Urban vegetation* as derived from the multispectral vegetation index NDVI at 30-m spatial resolution produced from Landsat TM (Figure 7)





Figure 7. Composite NDVI image of Athens for the warm season (April – August 2005).

3. *Land surface emissivity* (LSE) in the thermal infrared at 30-m spatial resolution produced from Landsat TM (Figure 8). The methodology applied is described in Sobrino et al. (2004) and Stathopoulou et al. (2007).






**Figure 8.** Seasonal surface emissivity image of Athens (April – August 2005).

Furthermore, processing of the satellite data in the near-infrared may lead to the measurement of the following biophysical variable for Athens:


4. **Land surface temperature (LST)** at 120-m spatial resolution produced from the Landsat TM sensor (Figure 9). The LST algorithm applied is the one proposed by Jiménez-Munóz and Sobrino (2003).



**Figure 9.** LST image of Athens on 10/7/2004.

### Data collected during the project

- *GIS data:* Vector maps for the Egaleo test site provided in shapefile format: census block, building block population, topography, land use, road network, bus routes, bus stops, education and sport units, healthcare, hydronet, rainnet, sewers, stream, manholes, monuments, squares, churches, points of interest, urban value zones, buildings (xls format), building heights (xls format) (see deliverable D3.3).
- *Satellite data*  
Three Landsat 5 images of the greater Athens area.
- *Aerial images*
- *Digital Elevation Model*  
Extraction and post processed ASTER GDEM for the broader area of Athens was provided by FORTH (Figure 10). The product was delivered in both Geographic WGS84 and UTM WGS84 projections with EGM96 vertical datum. Its spatial resolution is 30 m. A validation of this product has given by Chrysoulakis et al. (2009).



**Figure 10.** ASTER derived DEM for the broader area of Athens.

### Contact info

Dr. Marina Stathopoulou ([mstathop@phys.uoa.gr](mailto:mstathop@phys.uoa.gr))  
 Research Assistant, Remote Sensing and Image Processing Laboratory,  
 Department of Physics, University of Athens.  
 Tel.: +30 210 7276843



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 19/32

---



# BRIDGE

## Remote Sensing Data Collection and Analysis


Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 20/32

### 3.3 London

#### Data already existing

- **3D Maps:**  
Virtual London, Centre for Advanced Spatial Analysis (CASA), University College London (KCL: we can not provide the raw data)  
Data: may be able to be obtained via contact through Rachel Conti (Greater London Authority according to KCL)
  - **GIS data:**  
**SOTON GIS data**  
1 : 200 000 scale vector boundary data maps of Great Britain (Great Britain boundary data, Collins Bartholomew, UK), used to define urban features and London borough boundaries. The London Atmospheric Emissions Inventory (LAEI) for 2006 and forward projections (2015). This includes both the 25 m PM10 concentration map ( $\mu\text{g m}^{-3}$ ) and the total emission predictions for London (tonnes) for each borough and emission sources (1 km grid squares). A 25 m<sup>2</sup> resolution land classification map of the UK (LCM2000, Centre of Ecology and Hydrology, UK; Fuller et al., 2001)
  - **Satellite data.**
  - **Aerial images:**  
**KCL:** Airborne (one daytime, one night time flight in August 2008), north/south transects of Greater London:
    - hyperspectral radiances measured by NERC ARSF EAGLE/HAWK sensors, broadband radiances measured by NERC ARSF ATM (Spectral bands from 400 up to 2500 nm)
    - Leica ALS50-II LiDAR
    - Leica RCD105 39 megapixel digital camera
    - Accompanied by ground based measurements
- KCL: Airborne data collected June 2010 (one day time and one night flight) by NERC ARSF:
- hyperspectral radiances measured by Aisa EAGLE and HAWK sensors
  - thermal radiances measured by Itres TABI
  - Leica ALS50-II LiDAR
  - Leica RCD105 39 megapixel digital camera
  - Accompanied by ground based measurements e.g.
    - thermal imagers: FLIR SC640, MIRICLE K370, TESTO
    - surface spectrometers: GER1500, ASD Fieldspec3

**Figure 11.** Examples of surface characteristics at 1 km scale for London (a) built fraction (b) water fraction (c) mean building height



## Data to be collected during the project


- *GIS data*

- *SOTON GIS data*

Urban vegetation data for the Greater London Authority (GLA) has been extracted at each of the 33 London borough levels using ArcMap™ (ESRI® ArcMap™ version 9.3). The habitat classes extracted are: Class (1) Broad leaved trees, Class (2) Coniferous trees and Classes (5-8) Grasses.

The raster data-sets (25 m) were used to quantify the area of these land cover classes for each London borough.

Work has recently focused on the London Central Activity Zone and PM10 uptake to urban greenspace has been estimated using 20 m pixels as follows:



**Figure 12.** An uptake potential map of  $PM_{10}$  by the urban tree canopy. This potential uptake map shows the South West corner of the London CAZ and displays areas of existing broadleaf and coniferous woodland. For each 20 m pixel the estimated grams of  $PM_{10}$  uptake by annual dry deposition are given. The map was produced using three sources, the landcover 2000 map (Fuller et al., 2001), the  $PM_{10}$  dispersion map for the GLA (LAEI 2006) and  $PM_{10}$  deposition parameters from the UFORE model (Nowak, 1994).

### ***KCL GIS data***

Morphological characteristics

- Roughness length for momentum
- Zero plane displacement length
- Plan area index
- Frontal area index
- Sky view factors

Anthropogenic heat flux


[http://geography.kcl.ac.uk/micromet/wordpress/?page\\_id=436](http://geography.kcl.ac.uk/micromet/wordpress/?page_id=436)

Additional more detailed data are being determined for London  
 Surface Attributes for London: 1 km Grid scale

- further analysis of morphology, anthropogenic heat flux

RS& GIS & Meteorological data: turbulent sensible heat flux based on different modelling approaches (Aerodynamic Roughness Method, LUMPS)

- *Satellite data*  
**KCL:** Landsat Surface Temperature for London
- *Aerial images*  
**KCL:** 2008 Airborne observation will be repeated hopefully
- *Digital Elevation Model*  
Extraction and post processed ASTER GDEM for the broader area of London was provided by FORTH (Figure 11). The product was delivered in both Geographic WGS84 and UTM WGS84 projections with EGM96 vertical datum. Its spatial resolution is 30 m.



**Figure 13.** ASTER derived DEM for the broader area of London.

### Contact info

KCL:

Prof. Sue Grimmond ([sue.grimmond@kcl.ac.uk](mailto:sue.grimmond@kcl.ac.uk))

Dr Sean Beevers ([sean.beevers@kcl.ac.uk](mailto:sean.beevers@kcl.ac.uk))

SOTON:

Prof. Gail Taylor ([G.Taylor@soton.ac.uk](mailto:G.Taylor@soton.ac.uk))

Dr. M J. Tallis ([M.J.TALLIS@soton.ac.uk](mailto:M.J.TALLIS@soton.ac.uk))



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 24/32

### 3.4 Firenze

#### Data already existing

- *3D Maps*: Firenze 3D: <http://sit.comune.fi.it/volo.asp>  
Building height is a layer included in the CTR cartography described below
- *GIS data*  
DEM at 10m resolution if needed  
Alternatively, DEM at 90m resolution (SRTM)  
CORINE Land Cover at low resolution
- *Satellite data*
- *Aerial images*: aerial ortophoto if needed (2007? or 2003)
- *Other*  
Digital topographic maps at high resolution CTR (1:2000 scale), with layers describing roads /type etc), buildings type, (including height), green areas etc.

The products: Land Use/Land Cover data, Road Networks - Buildings, Topography, Population, Administrative units and more are now available for download at: <ftp://149.139.16.152>  
username: toscano, pwd: piero1980

A CODE\_LAYERS.txt for technical description (in english) is available. All layers are in shapefile format with attributes in italian language (e.g., quotagrond means base roof height).

#### Data collected during the project

- *Remote sensing data*  
Lidar PBL soundings (retrieving PBL characteristics + particulate profiles, ISAFOM), starting January 2010
- *GIS data*  
CMCC: Maps of sensible heat flux and surface emissivity will be produced using the WRF-ACASA model. High resolution maps (200 m) of surface emissivity, sensible and latent heat flux have been produced for the period January-April 2008 (Falk et al., 2010).
- *Satellite data*
- *Airborne acquired images*  
RS flight (IR + multispectral VIS-NIR) performed by ISAFOM
- *Digital Elevation Model*  
Extraction and post processed ASTER GDEM for the broader area of Firenze was provided by FORTH. The product was delivered in both Geographic WGS84 and UTM WGS84 projections with EGM96 vertical datum. Its spatial resolution is 30 m.

#### Contact info

CMCC:

Prof. Donatella Spano ([spano@uniss.it](mailto:spano@uniss.it))

Via de Nicola 9, 07100, Sassari, Italy

Tel. +39 079 229339

Dr. Serena Marras ([serenam@uniss.it](mailto:serenam@uniss.it))

Via de Nicola 9, 07100, Sassari, Italy

Tel. +39 079 229372

CNR-IBIMET:





# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 25/32

---

Beniamino Gioli ([b.gioli@ibimet.cnr.it](mailto:b.gioli@ibimet.cnr.it))  
Piero Toscano ([p.toscano@ibimet.cnr.it](mailto:p.toscano@ibimet.cnr.it))



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
Contract no.: 211345  
Document Ref.: 211345\_014\_TR\_CMCC  
Issue: 1.0  
Date: 18/05/2011  
Page number: 26/32

### 3.5 Gliwice

#### Data already existing

- *3D Maps*: no 3D maps available
- *GIS data*
  - About 30 000 GIS coverages in Gliwice Municipal Spatial Information System (MSIS). This coverages are available on 3 access level. First is for general user, the second is for investors (one has to register) and third, restricted for the town office clerk. One may view the first two on the homepage <http://test.um.gliwice.pl/portal/map>
  - Gliwice MSIS contains among others map of address points, map of buildings, lots, road network, utility networks, etc.
  - Gliwice MSIS also contains Digital Elevation Model in raster format. Resolution about 2 metres.
  - IETU has several meso-scale GIS coverages concerning environment and ecology including emission of PM10 in 500 metres grid.
- *Satellite data*: No satellite data available
- *Aerial images*: aerial orthophotomap serving as a background from year 2003

#### Data collected during the project

- *3D Maps*
- *GIS data*
- *Satellite data*
- *Airborne acquired images*
- *Digital Elevation Model*

Extraction and post processed ASTER GDEM for the broader area of Gliwice was provided by FORTH. The product was delivered in both Geographic WGS84 and UTM WGS84 projections with EGM96 vertical datum. Its spatial resolution is 30 m.

#### Contact info

IETU Katowice:  
Dr. Joachim Bronder ([bron@ietu.katowice.pl](mailto:bron@ietu.katowice.pl))  
ul. Kossutha 6, 40-844 Katowice  
Tel. +48 32 254 60 31

Gliwice Town Hall :  
Paweł Filipiak ([filipiak\\_p@sip.um.gliwice.pl](mailto:filipiak_p@sip.um.gliwice.pl))  
Tel.: 696 44 57 57



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
 Contract no.: 211345  
 Document Ref.: 211345\_014\_TR\_CMCC  
 Issue: 1.0  
 Date: 18/05/2011  
 Page number: 27/32

## Annex:

Remote Sensing measurements to be collected by each partner during the project. Methods, starting and closing date of data acquisition are also described.

**Table 2. FORTH.**

participant		Foundation for Research and Technology - Hellas					
acronym		FORTH					
case study		Athens					
Variable	type	Method	Equipment	starting date	closing date	frequency	Notes
Land surface temperature	Passive RS	Satellite remote sensing: Surface emissivity extraction, precipitable water and aerosols estimation, long-wave radiative transfer.	Software: ERDAS Imagine; ATCOR; SBDART. Data: Landsat and ASTER images for Athens	Jan 2010	Dec 2010		Spatial resolution: 60 - 120 m. It can be done for any of the case studies if the respective satellite data and ground truthing is available.
Land surface radiance, reflectance and brightness temperature	Passive RS						
Vegetation	Passive RS	Satellite remote sensing: Vegetation Indices; Fractional vegetation cover; Subpixel classification; Context-based classification.	Software: ERDAS Imagine; ENVI; Feature Analyst. Data: Landsat, SPOT, ASTER and Ikonos images for Athens	Jan 2010	Dec 2010		Spatial resolution: 1-30 m. It can be done for any of the case studies if the respective satellite data and ground truthing is available.
Topography	Passive RS	3D SURFACE CHARACTERISTICS. Satellite remote sensing: Photogrammetric analysis (DEM); GIS analysis (Slope, aspect, curvatures, watershed analysis).	Software: ERDAS Imagine; PCI Geomatics; SILCAST; ArcGIS. Data: ASTER images for Athens	Dec 2009	Dec 2009		Spatial resolution: 15 m. It can be done for any of the case studies if ASTER data are available.
Sensible Heat Flux	Active RS						
Surface Emissivity	Active RS						



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
 Contract no.: 211345  
 Document Ref.: 211345\_014\_TR\_CMCC  
 Issue: 1.0  
 Date: 18/05/2011  
 Page number: 28/32

**Table 3. KCL.**

	participant	King's College London United Kingdom					
	acronym	KCL					
	case study	London					
Variable	type	Method	Equipment	starting date	closing date	frequency	Notes
Land surface temperature	Passive RS	Satellite Remote Sensing	LandSAF land surface temperature (LST) product	all data relative to our observations will be analysed	Data download will continue after the end of the project	15 min	Spatial resolution: 60 - 120 m. It can be done for any of the case studies if the respective satellite data and ground truthing is available.
Land surface radiance, reflectance and brightness temperature	Passive RS	Satellite Remote Sensing; Airborne and LiDAR	Meteosat satellite data	RS flights dates are dependent on weather	LiDAR data will continue after the end of the project	15 min	
Vegetation	Passive RS						Spatial resolution: 1-30 m. It can be done for any of the case studies if the respective satellite data and ground truthing is available.
Topography	Passive RS						Spatial resolution: 15 m. It can be done for any of the case studies if ASTER data are available.
Sensible Heat Flux	Active RS	Satellite Remote Sensing, GIS and Meteorological data. Sensible heat flux calculation based on different modelling approaches (Aerodynamic Roughness Method, LUMPS)		all data relative to our observations will be analysed	Data download will continue after the end of the project		
Surface Emissivity	Active RS						



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
 Contract no.: 211345  
 Document Ref.: 211345\_014\_TR\_CMCC  
 Issue: 1.0  
 Date: 18/05/2011  
 Page number: 29/32

**Table 4. CNR.**

	participant	Consiglio Nazionale delle Ricerche					
	acronym	CNR					
	case study	Firenze					
Variable	type	Method	Equipment	starting date	closing date	frequency	Notes
Land surface temperature	Passive RS	airborne acquired images	FLIR M-40	giu-10		campaign	airborne acquired images
Land surface radiance, reflectance and brightness temperature	Passive RS	airborne acquired images	Geospatial Systems MS4100 CIR multispectral camera	giu-10		campaign	airborne acquired images
Vegetation	Passive RS						
Topography	Passive RS						
Sensible Heat Flux	Active RS						
Surface Emissivity	Active RS						



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
 Contract no.: 211345  
 Document Ref.: 211345\_014\_TR\_CMCC  
 Issue: 1.0  
 Date: 18/05/2011  
 Page number: 30/32

**Table 5. NKUA.**

	participant	National and Kapodistrian University of Athens					
	acronym	NKUA					
	case study	Athens					
Variable	type	Method	Equipment	starting date	closing date	frequency	Notes
Land surface temperature	Passive RS	Satellite Remote Sensing	Landsat TM/ETM+, ASTER LST	satellite images are acquired at periods or days that observation is needed	satellite images are acquired at periods or days that observation is needed	depends on the temporal resolution of the sensor: for TM, ETM+, ASTER: 16 days	
Land surface radiance, reflectance and brightness temperature	Passive RS	Satellite Remote Sensing	Landsat TM/ETM+, ASTER LST	satellite images are acquired at periods or days that observation is needed	satellite images are acquired at periods or days that observation is needed	depends on the temporal resolution of the sensor: for TM, ETM+, ASTER: 16 days	
Vegetation	Passive RS	Satellite Remote Sensing	Landsat TM/ETM+, ASTER LST	satellite images are acquired at periods or days that observation is needed	satellite images are acquired at periods or days that observation is needed	depends on the temporal resolution of the sensor: for TM, ETM+, ASTER: 16 days	Vegetation indices
Topography	Passive RS	surface slope, surface aspect, digital elevation model (DEM)	ASTER data	satellite images are acquired at periods or days that observation is needed	satellite images are acquired at periods or days that observation is needed	depends on the temporal resolution of the sensor: for TM, ETM+, ASTER: 16 days	
Sensible Heat Flux	Active RS						
Surface Emissivity	Active RS						



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
 Contract no.: 211345  
 Document Ref.: 211345\_014\_TR\_CMCC  
 Issue: 1.0  
 Date: 18/05/2011  
 Page number: 31/32

**Table 6. CMCC.**

	participant	Centro Euro-Mediterraneo per i Cambiamenti Climatici					
	acronym	CMCC					
	case study	Firenze					
Variable	type	Method	Equipment	starting date	closing date	frequency	Notes
Land surface temperature	Passive RS						
Land surface radiance, reflectance and brightness temperature	Passive RS						
Vegetation	Passive RS						
Topography	Passive RS						
Sensible Heat Flux	Active RS	modelling	ACASA model for flux simulation	set-09	giu-11		Initial conditions from in situ measurements; 30-min average; units: $W m^{-2}$
Surface Emissivity	Active RS	modelling	ACASA model for flux simulation	set-09	giu-11		Initial conditions from in situ measurements; 30-min average



# BRIDGE

## Remote Sensing Data Collection and Analysis

Deliverable no.: D.3.2.3  
 Contract no.: 211345  
 Document Ref.: 211345\_014\_TR\_CMCC  
 Issue: 1.0  
 Date: 18/05/2011  
 Page number: 32/32

**Table 7. SOTON.**

	participant	University of Southampton					
	acronym	SOTON					
	case study	London					
Variable	type	Method	Equipment	starting date	closing date	frequency	Notes
Land surface temperature	Passive RS						
Land surface radiance, reflectance and brightness temperature	Passive RS						
Vegetation	Passive RS	Use the following existing data sets: (i) Landcover 2000 (CEH), (ii) Trees in Towns II (Britt & Johnson 2008) and (iii) Chainsaw massacre (GLA, 2007) to estimate canopy cover type areas for GLA.		feb-09	ott-10		Estimated pollutant fluxes to the London Central activity Zone (Sept & October 2010).
Topography	Passive RS						
Sensible Heat Flux	Active RS						
Surface Emissivity	Active RS						